

**Serial No.: 09/998,058**

**IN THE CLAIMS:**

1. (Original) A method for identifying a genetic locus that modulates a phenotype, the method comprising:

- (a) providing a renewable population of genetically diverse individuals; and
- (b) mapping the genomes of individuals within the renewable population of genetically diverse individuals that display the phenotype, whereby a genetic locus that modulates the phenotype is identified.

2. (Original) The method of claim 1, wherein the renewable population of genetically diverse individuals comprises:

- (a) individuals produced by intercrossing recombinant inbred lines;
- (b) individuals produced by backcrossing recombinant inbred lines;
- (c) a cloned population of genetically diverse individuals; or
- (d) a panel of cell lines derived from genetically diverse individuals.

3. (Original) The method of claim 1, wherein an individual of the renewable population of genetically diverse individuals comprises a diploid, tetraploid, or polyploid organism, or a cell derived there from.

4. (Original) The method of claim 3, wherein the organism is selected from a group consisting of an animal and a plant.

5. (Original) The method of claim 4, wherein the animal is a mammal.

6. (Original) The method of claim 5, wherein the mammal is a rodent.

7. (Original) The method of claim 6, wherein the rodent is a mouse.

8. (Original) The method of claim 2, wherein the recombinant inbred lines comprise less than about 500 lines.

**Serial No.: 09/998,058**

9. (Original) The method of claim 8, wherein the recombinant inbred lines comprise less than about 100 lines.

10. (Original) The method of claim 2, wherein the recombinant inbred lines comprise one or more recombinant inbred lines selected from the group consisting of mouse lines AXB, BXA, CXB, and BXD.

11. (Original) The method of claim 2, wherein the recombinant inbred lines comprise recombinant inbred lines derived from at least 3 non-recombinant parent lines.

12. (Original) The method of claim 11, wherein the recombinant inbred lines comprise recombinant inbred lines derived from at least 4 non-recombinant parent lines.

13. (Original) The method of claim 12, wherein the recombinant inbred lines comprise recombinant inbred lines derived from at least 8 non-recombinant parent lines.

14. (Original) The method of claim 11, wherein the at least three non-recombinant parent lines comprise one or more non-recombinant parent lines selected from the group consisting of mouse lines C57BL/6, BALB/c, C3H, A, 129, and DBA/2.

15. (Original) The method of claim 2, wherein the cloned population or the panel of cell lines is derived from recombinant inbred line intercrosses, recombinant inbred line backcrosses, an F2 population, or a natural population.

16. (Original) The method of claim 15, wherein the recombinant inbred lines comprise recombinant inbred lines derived from at least 3 non-recombinant parent lines.

Serial No.: 09/998,058

17. (Original) The method of claim 16, wherein the recombinant inbred lines comprise recombinant inbred lines derived from at least 4 non-recombinant parent lines.

18. (Original) The method of claim 17, wherein the recombinant inbred lines comprise recombinant inbred lines derived from at least 8 non-recombinant parent lines.

19. (Original) The method of claim 1, wherein the mapping comprises analysis of genetic polymorphisms segregating in the renewable population of genetically diverse individuals.

20. (Original) The method of claim 1, wherein the phenotype is selected from the group consisting of a visible phenotype, a physiological phenotype, a behavioral phenotype, a susceptibility phenotype, a cellular phenotype, a molecular phenotype, and combinations thereof.

21. (Original) The method of claim 20, wherein the molecular phenotype is selected from the group consisting of a level of gene expression, a splice selection, a level of protein, a protein type, a protein modification, a level of lipid, a lipid type, a lipid modification, a level of carbohydrate, a carbohydrate type, a carbohydrate modification, and combinations thereof.

22. (Original) The method of claim 1, wherein the phenotype is modulated by a non-genetic factor.

23. (Original) The method of claim 22, wherein the phenotype is modulated by an interaction between two or more non-genetic factors.

24. (Original) The method of claim 22, wherein the non-genetic factor is an environmental condition or drug exposure.

Serial No.: 09/998,058

25. (Original) The method of claim 1, wherein the phenotype is modulated by an interaction between a genetic locus and a non-genetic factor.

26. (Original) The method of claim 25, wherein the non-genetic factor is an environmental condition or drug exposure.

27. (Original) The method of claim 1, further comprising identifying two or more genetic loci that modulate the phenotype.

28-45. (Currently Canceled)

46. (Original) A method for identifying an interaction between a genetic locus and a non-genetic factor, wherein the interaction modulates a phenotype, the method comprising:

- (a) providing a renewable population of genetically diverse individuals;
- (b) providing a non-genetic factor to the renewable population; and
- (c) mapping the genomes of individuals that display the phenotype, whereby an interaction between a genetic locus and the non-genetic factor that modulates a phenotype is identified.

47. (Original) The method of claim 46, wherein the renewable population of genetically diverse individuals comprises:

- (a) individuals produced by intercrossing recombinant inbred lines;
- (b) individuals produced by backcrossing recombinant inbred lines;
- (c) a cloned population of genetically diverse individuals; or
- (d) a panel of cell lines derived from genetically diverse individuals.

48. (Original) The method of claim 46, wherein the phenotype is selected from the group consisting of a visible phenotype, a physiological phenotype, a behavioral

Serial No.: 09/998,058

phenotype, a susceptibility phenotype, a cellular phenotype, a molecular phenotype, and combinations thereof.

49. (Original) The method of claim 48, wherein the molecular phenotype is selected from the group consisting of a level of gene expression, a splice selection, a level of protein, a protein type, a protein modification, a level of lipid, a lipid type, a lipid modification, a level of carbohydrate, a carbohydrate type, a carbohydrate modification, and combinations thereof.

50. (Original) The method of claim 46, further comprising identifying an interaction among two or more genetic loci and a non-genetic factor.

51. (Original) The method of claim 46, further comprising identifying an interaction among a genetic locus and two or more non-genetic factors.

52. (Original) The method of claim 46, wherein the non-genetic factor is an environmental condition or drug exposure.

53. (Original) The method of claim 46, further comprising identifying an interaction among two or more genetic loci and two or more non-genetic factors.

54-59. (Currently Canceled)